Review of Contemporary Philosophy
ISSN: 1841-5261, e-ISSN: 2471-089X
Vol 22 (1), 2023
Pp 4669 - 4683

# The Role of Pharmacists in Supporting Nurses to Improve Medication Adherence in Chronic Disease Patients

**[1]hussein Abdullah Mohammad Almahdi,[2]. Faisal Salah Alharbi,[3]. Alyah Sadiq Mohammed Alismail,[4]. Duaa Sadiq Nasser Albeladi,[5]. Hussain Ali Al Bahrani,[6]. Fatimah Ahmed Tuhaifah,[7]. Fatimah Habib Alsaeed,[8] Abeer Hamad Al-Hejili,[9]. Ayat Saleh Al Thani,[10] Zainab Ali Alnajim,[11]. Layla Abdullah Al Hussain,[12]. Ahood Nughaymish Almutiari,[13]. Bader Nughaymish Almutairi,[14]. Narjes Ali Alyagoub,[15]- Fatimah Radhi Almahasnah**

[1] PHARMACY MURITBAH PHC

[2] Pharmacy Technician Al-Rass General Hospital

[3]technician Pharmacy Dammam Medical Complex

[4]pharmacy Technician Prince Sultan Cardiac Center

[5] Clinical Pharmacist Prince Saud Bin Jalawi Hospital

[6] Pharmacy Technician Dammam Medical Complex

[7]paramedic Ministry Of Health Helth Emergency Operations Centre

[8]nursing Technician Bab Al Majidi Health Center

[9] Nursing Technician Safwa General Hospital

[10] Nursing Technician Public Health Administration - School Health

[11]nursing Technician Anak General Hospital

[12] Nursing Al-Qaisumah Hospital , Health Cluster,

[13]nursing Hafar Al-Batin Central Hospital, Health Cluster,

[14] Nursing (Nqps) Nursing Quality And Patient Safety: Aljubail General Hospital

[15] Nursing Specialist Al-Qadisiyah Health Center Dammam, Ksa

**Received:** 16 october 2023 **Revised:** 29 November 2023 **Accepted:** 13 December 2023

## Chapter 1: Introduction to Medication Adherence in Chronic Disease Management

Medication adherence is a crucial factor in the effective management of chronic diseases such as diabetes, hypertension, and cardiovascular diseases. Chronic conditions often require long-term or lifelong treatment regimens, where consistent medication use is essential to control symptoms, prevent complications, and improve quality of life. When patients adhere to prescribed treatments, the likelihood of disease progression is significantly reduced, leading to better health outcomes and reduced hospitalizations **(Kvarnström et al., 2021).** Conversely, poor adherence can result in treatment failures, exacerbations of symptoms, and increased healthcare costs. Studies consistently show that medication non-adherence is a significant contributor to the global burden of chronic diseases. Therefore, ensuring that patients follow their treatment regimens is a fundamental challenge in healthcare, requiring tailored interventions and the active involvement of various healthcare professionals, including nurses and pharmacists **(Sugandh et al., 2023).**

Patients with chronic diseases face numerous barriers to medication adherence, which can be categorized into factors related to the patient, the medication itself, and the healthcare system. Patient-related barriers include forgetfulness, lack of understanding about the importance of the medication, and the emotional burden of managing a long-term condition. Medication-related barriers may include side effects, complex dosing schedules, and the cost of treatment, which can be prohibitive for some patients **(Konstantinou et al., 2020).** Healthcare-related challenges include poor communication with healthcare providers, lack of follow-up, and limited access to resources. The interplay of these factors often leads to inconsistent medication use, which compromises treatment efficacy. Addressing these barriers requires a comprehensive, patient-centered approach that considers both the physical and psychological aspects of managing a chronic condition **(Kelley et al., 2022).**

The role of healthcare professionals in improving medication adherence is critical. Nurses, pharmacists, and physicians each bring unique skills and perspectives to the management of chronic diseases. Pharmacists provide expertise in drug therapy management, ensuring that patients receive the right medication at the correct dose, while nurses are integral in offering continuous education, emotional support, and monitoring of treatment outcomes. Collaborative efforts between healthcare providers increase the likelihood of identifying and addressing barriers to adherence early on **(De Baetselier et al., 2021).** A multidisciplinary approach can significantly enhance patient understanding, motivation, and confidence in their treatment plans. By working together, pharmacists and nurses can identify and resolve issues such as misunderstanding medication instructions, difficulty with administration, or financial constraints, thereby improving adherence and patient outcomes **(Alemede et al., 2024).**

Poor medication adherence has a profound impact on both patients and the healthcare system. Non-adherence is associated with an increased risk of hospitalization, disease complications, and even death. For example, patients with uncontrolled hypertension may suffer from strokes or heart attacks due to inconsistent medication use **(Gil-Guillen et al., 2022).** Similarly, diabetes patients who do not take their insulin as prescribed may experience complications such as neuropathy, kidney failure, or blindness. Beyond health consequences, non-adherence also leads to higher healthcare costs due to avoidable emergency room visits, hospitalizations, and additional treatments. The financial burden on both patients and healthcare systems underscores the importance of strategies to improve adherence. Therefore, interventions that increase medication adherence are essential not only for improving individual patient outcomes but also for reducing the overall cost of chronic disease management **(Burnier, 2024).**

Medication adherence is a multifaceted process that involves patient behavior, education, and communication with healthcare providers. It is not simply about taking medications on time; it is about understanding the importance of the treatment, managing side effects, and maintaining the motivation to continue treatment over time. For patients with chronic diseases, adherence is often a lifelong commitment. Education about the purpose of the medication, potential side effects, and the consequences of non-adherence can empower patients to take an active role in managing their health **(Pruette & Amaral, 2021).** Additionally, fostering a trusting relationship between patients and healthcare providers, particularly nurses and pharmacists, can encourage open communication, allowing patients to voice concerns and receive support. Addressing both the psychological and practical aspects of adherence is essential in improving outcomes **(Alsaihati et al., 2024).**

Pharmacists are key members of the healthcare team when it comes to improving medication adherence. As medication experts, pharmacists are well-positioned to offer personalized counseling on how to properly use medications, explain side effects, and monitor for potential drug interactions. They can assist in simplifying medication regimens, such as by recommending fixed-dose combination medications to reduce pill burden, or providing medication synchronization services to align refill dates **(Balogun et al., 2024).** Pharmacists can also engage in medication therapy management (MTM) programs to ensure that patients are following their prescribed treatments appropriately. These programs often involve a comprehensive review of the patient's medication list and counseling on optimizing medication use. By working alongside nurses, pharmacists can help reinforce the importance of adherence and address any medication-related concerns or challenges patients might have **(Ferreri et al., 2020).**

Nurses are often the most frequent point of contact for patients with chronic diseases, making them crucial in supporting medication adherence. Nurses play an essential role in educating patients about their conditions and treatments, reinforcing the information provided by pharmacists and doctors. They can offer practical advice on medication management, such as how to incorporate medications into daily routines, and address concerns about side effects **(Yang et al., 2022).** Nurses can also monitor adherence by tracking medication usage during follow-up visits and identifying patients who may be struggling with non-adherence. They are uniquely positioned to provide emotional support, helping patients navigate the challenges of managing chronic conditions. Additionally, nurses can communicate any concerns or barriers identified by the patient to the rest of the healthcare team, ensuring a holistic approach to improving medication adherence **(While, 2020).**

A collaborative care model, where pharmacists and nurses work together to support medication adherence, has shown great promise in improving patient outcomes. This team-based approach ensures that patients receive consistent, comprehensive care throughout their treatment journey. Pharmacists and nurses can collaborate on patient education, share information about potential side effects, and work together to address barriers such as medication costs or complex dosing schedules **(Al-Worafi, 2023).** For example, nurses can identify patients who need more detailed explanations of their medication regimen, and pharmacists can provide the necessary support by offering counseling and alternative medication options. By working closely together, pharmacists and nurses can empower patients to take control of their health, increasing medication adherence and improving long-term outcomes in chronic disease management **(Alyami et al., 2024).**

A patient-centered approach is fundamental to improving medication adherence. This approach involves considering the individual needs, preferences, and challenges of each patient when developing a treatment plan. Nurses and pharmacists must take the time to understand the patient's lifestyle, health goals, and any barriers they may face in adhering to their medication regimen **(Hamrahian et al., 2022).** For example, a patient with diabetes may struggle with remembering to take their medication due to a busy schedule or difficulty with understanding instructions. A patient-centered approach would involve simplifying the regimen, providing medication reminders, and offering education that aligns with the patient’s understanding and capabilities. Nurses and pharmacists must also be empathetic, recognizing the emotional and psychological factors that may impact adherence, such as fear of side effects or denial of the severity of the disease. This personalized approach increases patient engagement and promotes better adherence **(Oakley & Ream, 2024).**

Advancements in technology have introduced new tools and solutions to improve medication adherence. Digital health tools, such as medication reminder apps, smart pill bottles, and electronic health records (EHRs), allow both nurses and pharmacists to monitor adherence and engage patients in their care. These tools can send reminders to patients to take their medications on time, track their progress, and even alert healthcare providers if a patient misses doses **(Toscos et al., 2020).** EHRs enable better communication between healthcare teams, ensuring that nurses and pharmacists are aware of any changes in the patient’s medication regimen. Telemedicine services also allow for remote consultations, where pharmacists and nurses can check in with patients to discuss their medications and adherence levels. By leveraging technology, healthcare professionals can provide timely support and interventions, ultimately enhancing adherence and improving health outcomes for patients with chronic diseases **(Haleem et al., 2021).**

Cultural and social factors play a significant role in medication adherence, especially in patients from diverse backgrounds. Language barriers, cultural beliefs about illness and medication, and socioeconomic factors can all impact a patient’s willingness or ability to follow their prescribed treatment plan **(Kasahun et al., 2022).** For example, some patients may prefer traditional or alternative therapies over conventional medications due to cultural practices. Nurses and pharmacists must be sensitive to these factors and adapt their approach accordingly. This might include offering educational materials in the patient’s preferred language, respecting cultural beliefs, and helping patients access affordable medications. Building trust with patients and understanding their cultural context is essential in overcoming these barriers and improving adherence. Addressing the social determinants of health, such as income and access to healthcare, also plays a crucial role in ensuring that patients can follow their medication regimen consistently **(Seixas et al., 2020).**

The future of medication adherence in chronic disease management lies in a more personalized, technology-driven approach that integrates healthcare professionals' expertise with patients' individual needs. With the growing use of artificial intelligence, machine learning, and personalized medicine, healthcare providers will be able to offer more tailored treatment regimens and identify adherence challenges earlier. Pharmacists and nurses will continue to play an essential role in this process, working together to provide more efficient and patient-centered care **(Coman et al., 2024).** By incorporating digital tools, enhancing communication, and addressing individual barriers to adherence, healthcare teams can ensure better management of chronic diseases, reduce hospitalizations, and improve the overall quality of life for patients. The goal is not only to ensure that patients take their medications but also to empower them to actively participate in their healthcare journey, ultimately leading to better health outcomes and reduced healthcare costs **(Al-Jaroodi et al., 2020).**

Chapter 2: **Pharmacists' and Nurses' Roles in Medication Management and Patient Education**

Pharmacists play a critical role in medication management, ensuring that patients receive the correct medications in the right dosages, while minimizing the risk of adverse reactions. Pharmacists are experts in pharmacology, understanding how drugs interact with each other and the body. They are responsible for reviewing prescriptions, ensuring that there are no potential drug-drug or drug-disease interactions, and that the prescribed medications are appropriate for the patient's condition **(Al-Dosari et al., 2024).** By carefully evaluating a patient's medical history, current medications, and lab results, pharmacists can help prevent medication errors, ensuring that patients adhere to their treatment regimens effectively. This active involvement in medication management ensures that patients not only receive appropriate medications but also benefit from therapy that is tailored to their specific needs and circumstances **(Coutsouvelis et al., 2020).**

Medication Therapy Management (MTM) is a structured service provided by pharmacists to optimize therapeutic outcomes for patients. MTM involves comprehensive medication reviews, patient counseling, and ongoing monitoring. The goal of MTM is to ensure that the medications a patient is taking are effective, safe, and being used correctly **(Ferreri et al., 2020).** Pharmacists conducting MTM services will assess a patient's entire medication regimen, identifying any issues such as drug interactions, duplicate therapies, or medications that the patient may not be taking correctly. This service enhances patient understanding of their medications and can reduce the likelihood of medication non-adherence. By working with patients to simplify and improve their medication regimens, pharmacists help prevent complications and improve overall health outcomes, thus playing a crucial role in chronic disease management **(Jarab et al., 2022).**

Patient education is a fundamental aspect of pharmacy practice, as it empowers patients to take control of their health and treatment. Pharmacists provide education to patients on the correct use of medications, including when and how to take them, potential side effects, and what to do in case of a missed dose. They also educate patients about the importance of adherence to prescribed regimens, helping them understand that skipping doses or stopping medications prematurely can lead to worsened health conditions **(Alqarni et al., 2023).** Pharmacists are well-positioned to offer this education due to their expertise in pharmacology and the accessibility they have to patients. This knowledge can be particularly valuable in educating patients about chronic disease management, where long-term adherence is crucial to preventing complications and improving the quality of life **(Alemede et al., 2024).**

A key aspect of patient education is counseling on medication adherence. Pharmacists educate patients about the potential consequences of poor adherence, such as disease progression, hospitalization, or the development of complications. They can also provide practical strategies to help patients remember to take their medications, such as setting reminders, using pill organizers, or syncing medication schedules with daily routines **(Konstantinou et al., 2020).** Pharmacists can work with patients to identify barriers to adherence, such as difficulty swallowing pills, financial concerns, or complex medication regimens, and suggest alternatives. By addressing these challenges and offering personalized solutions, pharmacists can significantly increase the likelihood that patients will follow their prescribed treatment plans and achieve better health outcomes **(Balogun et al., 2024).**

Pharmacists play a key role in simplifying medication regimens to improve adherence. Many patients with chronic diseases are prescribed multiple medications, which can be overwhelming and difficult to manage **(Al-Anzi et al., 2024).** Pharmacists can review a patient's medication list and identify opportunities for simplification, such as switching to combination pills, adjusting dosing schedules, or recommending alternative formulations like liquid medications for patients who have difficulty swallowing pills. Simplifying a patient's medication regimen reduces the complexity and potential for mistakes, making it easier for patients to adhere to their treatment plan. This process not only improves adherence but also enhances patient satisfaction and engagement with their treatment, as patients are more likely to stay on track with a regimen that is manageable **(Darlow et al., 2023).**

Pharmacists are uniquely positioned to address patient concerns about medication side effects. Many patients worry about potential adverse effects when starting a new medication, which can sometimes lead to non-adherence. Pharmacists can provide clear, evidence-based information about the expected side effects of medications and help patients understand that some side effects are temporary or manageable **(Alemede et al., 2024)**. By explaining how to mitigate side effects, such as taking medications with food or adjusting the timing of doses, pharmacists can help patients feel more comfortable with their treatment regimens. Additionally, when patients experience serious side effects, pharmacists can act as a liaison between patients and physicians to adjust or change the treatment plan, ensuring that the patient remains adherent to a medication regimen that is both effective and well-tolerated **(Chavan et al., 2023).**

Chronic disease management relies heavily on medication adherence. Pharmacists educate patients about the importance of staying on their medication regimens to manage their conditions effectively. For patients with diseases such as diabetes, hypertension, or heart disease, consistent medication adherence can prevent complications and improve long-term health outcomes. Pharmacists can explain how each medication works to control symptoms, prevent disease progression, and reduce the risk of exacerbations or hospitalizations **(Hamrick et al., 2020).** By helping patients understand the direct connection between adherence and better health outcomes, pharmacists increase the likelihood that patients will see the value in maintaining their treatment plans, even when they may not feel immediate effects from their medications **(Abdulhussein et al., 2024)**.

Collaboration between pharmacists and nurses is essential for improving medication adherence in chronic disease patients. Nurses are often the first healthcare professionals to interact with patients and have a unique opportunity to reinforce the medication education provided by pharmacists **(Kiles et al., 2022).** By working together, pharmacists and nurses can create a unified approach to patient education, ensuring consistency in the messaging and addressing any questions or concerns patients may have. Pharmacists can provide nurses with the latest drug information, helping them stay informed about new medications or updated treatment guidelines, while nurses can identify any adherence issues they observe during patient interactions. Together, pharmacists and nurses can address barriers to adherence, such as misunderstandings about the medication regimen, by offering tailored solutions that meet the patient's individual needs **(Raza et al., 2022).**

Advancements in technology have provided pharmacists with new tools to support medication management and improve adherence. Electronic health records (EHRs) allow pharmacists to track patients' medication histories, identify potential issues, and intervene when necessary **(Ravi et al., 2022).** Pharmacists can also leverage mobile health applications to provide patients with reminders to take their medications, track their progress, and offer educational resources. These technologies also allow for real-time communication between pharmacists, nurses, and patients, ensuring that any changes in the treatment plan are promptly addressed. By integrating technology into medication management, pharmacists can enhance patient engagement and improve adherence rates, particularly for patients with chronic conditions who require long-term medication management **(Kiles et al., 2022).**

Pharmacists play a critical role in identifying and overcoming barriers to medication adherence. Financial constraints, lack of understanding, forgetfulness, and fear of side effects are common challenges that prevent patients from following their prescribed treatment plans. Pharmacists can offer practical solutions to these barriers, such as recommending generic medications to reduce costs, providing clear and simple explanations about the benefits of the treatment, and suggesting strategies to improve medication-taking habits **(Bethishou, 2024).** For example, pharmacists can help patients set up medication synchronization programs, where all medications are filled on the same day, simplifying the process. By addressing these barriers and offering patient-centered solutions, pharmacists can significantly improve adherence and help patients achieve better health outcomes **(Ngamdee et al., 2024).**

Pharmacists also play an ongoing role in monitoring medication adherence and providing feedback to patients and healthcare teams. By reviewing prescription refill records, conducting medication reviews, and interacting with patients, pharmacists can identify patients who may not be adhering to their medication regimens. For example, a delay in refills may signal a problem with adherence **(Ekong et al., 2020).** When these issues are detected, pharmacists can provide feedback to patients, reinforcing the importance of staying on track with their treatment plan. They can also work closely with nurses to discuss adherence issues and determine the most appropriate interventions. By continuously monitoring adherence and providing timely feedback, pharmacists help ensure that patients remain on track with their medication regimens, ultimately improving health outcomes **(Bethishou, 2024).**

The role of pharmacists in medication management and patient education is crucial for improving medication adherence, particularly in patients with chronic diseases. By leveraging their expertise in pharmacology, drug interactions, and patient counseling, pharmacists can ensure that patients understand the importance of adherence and can navigate the challenges they may face **(Patton et al., 2021).** When pharmacists collaborate with nurses, they can create a holistic support system that enhances patient education, simplifies medication regimens, and addresses barriers to adherence. Through these efforts, pharmacists contribute significantly to improving patient outcomes and enhancing the quality of care in chronic disease management. The integrated approach, combining the strengths of both professions, results in better health outcomes, reduced complications, and a more comprehensive care plan for patients **(Syrnyk & Glass, 2023).**

## Chapter 3: **Collaborative Strategies: How Pharmacists and Nurses Can Work Together to Improve Adherence**

Effective communication between pharmacists and nurses is a cornerstone of improving medication adherence in chronic disease patients. Clear and consistent communication helps ensure that both healthcare professionals are on the same page regarding the patient's treatment plan. Nurses often have firsthand knowledge of the patient's challenges with adherence, such as difficulty understanding instructions or dealing with side effects, and can share this with pharmacists **(White et al., 2022).** Pharmacists, in turn, can provide valuable insight into the pharmacological aspects of the treatment, suggesting alternatives or adjustments that might improve adherence. Regular discussions about the patient's progress allow for timely interventions, and joint decision-making fosters a sense of shared responsibility. Establishing clear communication protocols, such as regular team meetings or a shared electronic health record system, can help streamline this process and ensure that both pharmacists and nurses are actively involved in managing the patient’s medication regimen **(Ravi et al., 2022).**

When pharmacists and nurses align their goals and work together towards a common objective—improving medication adherence—they create a unified approach that benefits the patient. Nurses are often tasked with monitoring the patient’s behavior and providing support, while pharmacists can offer expert guidance on drug efficacy and potential side effects **(Schmid et al., 2022).** By setting measurable goals, such as a specific adherence rate or reduced frequency of medication errors, both professionals can assess the patient’s progress over time. These shared goals also help identify areas where additional support is needed. For example, if a patient is not meeting adherence goals, the team can jointly discuss strategies such as simplifying the medication regimen or addressing financial barriers. Having unified goals promotes collaboration, ensures that both professionals work towards the same outcomes, and ultimately improves the patient's health outcomes by increasing adherence **(Wheat et al., 2020).**

One of the most effective collaborative strategies is the development of joint patient education programs that involve both pharmacists and nurses. While pharmacists focus on educating patients about the technical aspects of their medications, such as dosage, side effects, and interactions, nurses can provide additional context regarding the patient's lifestyle and how the medication fits into their daily routine. Joint education sessions can take place in person, through group education programs, or via digital platforms **(Balogun et al., 2023)**. During these sessions, both professionals can reinforce key messages, ensuring that patients receive consistent information. For example, pharmacists might explain how a certain medication works, while nurses could discuss how to manage side effects in a real-world setting. By working together in education, pharmacists and nurses ensure that patients receive holistic, patient-centered care that is more likely to lead to improved medication adherence **(Dilles et al., 2021)**.

.Collaboration between pharmacists and nurses also facilitates the adoption of patient-centered care approaches, which are essential for improving medication adherence. Both professionals must understand the patient's individual needs, preferences, and challenges when it comes to medication management. This collaborative approach involves tailoring care plans to fit the patient's unique lifestyle and circumstances **(White et al., 2022)**. Nurses often have the best insight into the patient's daily routine, social circumstances, and potential barriers to adherence, such as forgetfulness or financial strain. Pharmacists, with their in-depth knowledge of pharmacology, can help adjust the treatment regimen to make it more manageable for the patient. For instance, a pharmacist might recommend once-daily medications or simpler regimens, while a nurse can provide reminders or offer techniques to incorporate medication adherence into the patient's routine. This personalized, team-based approach helps ensure the patient's success in adhering to the treatment plan **(Jaffer et al., 2023).**

Coordinating follow-up care is another critical strategy in improving medication adherence. Regular follow-ups with both pharmacists and nurses allow for continuous monitoring of the patient's progress and any issues related to medication adherence. Nurses, who often have more frequent contact with patients, can be the first to identify when adherence issues arise. They can alert the pharmacist to any potential problems, such as missed doses or adverse reactions, allowing the pharmacist to provide timely interventions **(Lott et al., 2021).** This collaboration helps prevent medication non-adherence from escalating and ensures that corrective actions are implemented quickly. Pharmacists can also offer counseling during follow-up appointments, reinforcing the importance of adherence and addressing any questions or concerns the patient might have. Coordinated follow-up care creates a system of ongoing support that improves the likelihood of long-term medication adherence **(Forsyth et al., 2023).**

Technology can play a significant role in enhancing collaboration between pharmacists and nurses in improving medication adherence. Digital health tools such as mobile apps, electronic health records (EHRs), and telehealth platforms enable real-time communication and coordination between the two professionals. For instance, EHR systems allow pharmacists and nurses to share patient information, track medication regimens, and monitor adherence patterns **(Coutsouvelis et al., 2020).** Mobile apps can send reminders to patients about when to take their medications, reducing the burden on nurses. These tools can also allow pharmacists to provide virtual consultations, offering patients easy access to expert advice without the need to visit the clinic. By leveraging technology, pharmacists and nurses can ensure that adherence support is accessible, timely, and personalized, making it easier for patients to follow their prescribed treatments **(Rahayu et al., 2021)**.

Pharmacists and nurses must work together to identify and address the common barriers that prevent patients from adhering to their medication regimens. Financial constraints, complex medication schedules, side effects, and lack of understanding about the importance of medications can all contribute to non-adherence. Nurses, with their close relationship with patients, are often the first to identify these barriers and can bring them to the attention of the pharmacist **(Al-Worafi, 2023).** Pharmacists, in turn, can recommend alternatives such as generics, dose simplifications, or medication therapy management programs to address these challenges. By collaborating on the management of these barriers, nurses and pharmacists can ensure that patients receive the support they need to overcome obstacles to adherence. Together, they can develop tailored strategies to address each patient's unique circumstances, ensuring a more personalized approach to medication management **(Lott et al., 2021).**

An important strategy for improving medication adherence is encouraging patient empowerment. Both pharmacists and nurses play key roles in helping patients understand their medications and take responsibility for their treatment **(Pruette & Amaral, 2021).** Nurses can educate patients about the importance of their medication in managing their chronic condition, while pharmacists can provide more detailed information about each drug's function, side effects, and potential interactions. Empowering patients through education builds their confidence and encourages active participation in their treatment. When patients feel more informed, they are more likely to make informed decisions about their medication and adhere to the prescribed regimen. Joint efforts between pharmacists and nurses to educate and empower patients lead to increased engagement and long-term medication adherence, improving patient outcomes and quality of life **(Schmid et al., 2022).**

When patients struggle with medication adherence, a collaborative problem-solving approach between pharmacists and nurses can help find viable solutions. Nurses may notice early signs of non-adherence, such as missed doses, and bring them to the attention of the pharmacist **(Tadros et al., 2023).** The pharmacist can then assess the situation and suggest potential solutions, such as switching to an easier-to-take medication, offering patient education on the importance of adherence, or providing access to financial assistance programs. This proactive approach allows for a tailored intervention that addresses the root causes of non-adherence. For instance, if a patient has difficulty swallowing pills, the pharmacist may suggest liquid alternatives or discuss pill-splitting techniques, while the nurse can work with the patient to ensure they understand the new regimen. By collaborating to solve adherence challenges, both professionals can enhance patient outcomes **(Passey et al., 2021).**

Organizing multidisciplinary team meetings, where pharmacists, nurses, and other healthcare providers come together to discuss patient care plans, can greatly enhance medication adherence. These meetings foster an environment of shared responsibility, where each team member brings their expertise to address challenges and opportunities for improvement. Nurses can provide valuable insights into the patient's behavior, lifestyle, and adherence patterns, while pharmacists can contribute their knowledge of medications and potential therapeutic adjustments **(O'Toole et al., 2022)**. Such meetings also help in creating a comprehensive care plan that includes strategies for improving adherence. For example, a nurse may suggest additional support or follow-up for a patient struggling with side effects, while the pharmacist can review the medication regimen to identify any potential adjustments that could improve adherence. Collaborative team meetings ensure that the entire care team is aligned in supporting the patient's medication adherence **(Balogun et al., 2023).**

Collaboration between pharmacists and nurses is most effective when patients trust both professionals. Building a strong rapport with patients is key to encouraging open communication and improving medication adherence. Nurses, who often have more frequent and personal interactions with patients, are in a prime position to establish this trust. Through compassionate care and active listening, nurses can create a safe environment where patients feel comfortable discussing their challenges with medication adherence **(Bhattarai et al., 2021)**. Pharmacists can further support this relationship by providing clear, understandable medication information and offering ongoing counseling. When both pharmacists and nurses work together to build trust with patients, they can identify underlying issues that may be hindering adherence, such as misconceptions about medications or fear of side effects. Trust enhances the effectiveness of collaboration and ultimately leads to improved patient outcomes **(Schneider & Burnier, 2023).**

### 3.12 Measuring and Evaluating Adherence Outcomes

Finally, measuring and evaluating the outcomes of collaborative efforts is essential for assessing the effectiveness of strategies to improve medication adherence. Both pharmacists and nurses can use various methods to track adherence, such as pill counts, medication refill rates, or self-reported adherence surveys. Regular evaluation allows for adjustments to be made to care plans if necessary, ensuring that interventions remain effective **(Jaffer et al., 2023).** In addition, documenting the results of these efforts can provide valuable feedback for both professionals, allowing them to refine their collaborative strategies for future patients. By monitoring the success of their interventions, pharmacists and nurses can ensure that they are continuously improving their approach to medication adherence, leading to better health outcomes for patients with chronic diseases. This process of evaluation and adaptation is a key component of providing high-quality, patient-centered care **(Tadros et al., 2023).**

## Chapter 4: **Barriers to Medication Adherence and Pharmacists' Role in Overcoming Challenges**

Medication adherence remains a significant challenge in managing chronic diseases. Patients often face multiple barriers that hinder their ability to take medications as prescribed. One of the most common barriers is the fear or experience of side effects. Many patients may stop taking their medications due to concerns about adverse reactions, even if these side effects are minor or temporary. Furthermore, patients with chronic conditions often require long-term treatment, which can lead to medication fatigue, causing them to overlook or discontinue their prescribed regimen **(Sugandh et al., 2023)**. Pharmacists can address these barriers by counseling patients on the expected side effects, discussing ways to manage them, and reassuring them about the benefits of continuing treatment despite minor discomfort. By providing this education, pharmacists help patients feel more comfortable with their medications, ultimately improving adherence **(Jarab et al., 2022).**

Cost is another significant barrier to medication adherence, particularly in patients with chronic diseases who may require multiple medications over an extended period. High out-of-pocket costs can lead patients to skip doses, reduce their medication regimen, or even stop treatment altogether **(Al-Jaroodi et al., 2020)**. Pharmacists play a crucial role in mitigating this challenge by helping patients find more affordable options. This could include recommending generic medications, exploring patient assistance programs, or advising on over-the-counter alternatives where appropriate. Pharmacists can also work closely with healthcare providers to adjust treatment plans that meet both the clinical needs and financial constraints of the patient. By addressing the cost factor, pharmacists enable patients to continue their treatment without the burden of financial stress, improving their long-term adherence and health outcomes **(Wheat et al., 2020).**

Forgetfulness is another common barrier to medication adherence, particularly for patients managing multiple chronic conditions. Daily regimens can be difficult to remember, especially if patients are managing several medications with different dosages or schedules. This often leads to missed doses, which can impact the effectiveness of treatment. Pharmacists can assist by suggesting strategies to improve medication tracking, such as using pill organizers, setting alarms, or using medication reminder apps **(Kvarnström et al., 2021)**. Additionally, pharmacists can encourage patients to involve family members or caregivers in medication management, ensuring that someone else can help remind the patient when it is time to take their medication. Pharmacists can also offer guidance on simplifying medication regimens by working with the healthcare team to reduce the number of medications or adjust schedules to make them easier for the patient to follow **(O'Toole et al., 2022).**

Many patients do not fully understand the importance of their medications, leading them to neglect or discontinue treatment. This lack of understanding may arise from unclear communication with healthcare providers or insufficient patient education. Pharmacists are essential in bridging this gap by ensuring patients understand why each medication is necessary and how it fits into their overall treatment plan **(Alqarni et al., 2023).** Pharmacists can explain the mechanism of action of each medication, how it helps manage the disease, and the potential consequences of non-adherence. This knowledge empowers patients to take ownership of their health and understand the long-term benefits of sticking to their prescribed regimen. Educating patients about the role of medications in controlling their condition, improving their quality of life, and preventing complications encourages adherence and helps establish trust in the treatment plan **(Bhattarai et al., 2021).**

Psychological factors, such as depression, anxiety, or fear, can significantly impact medication adherence. Patients experiencing mental health challenges may feel overwhelmed by their chronic condition and become disengaged from their treatment regimen. They may forget to take medications, or worse, refuse to take them due to feelings of hopelessness or helplessness **(Yang et al., 2022).** Pharmacists can play a critical role in identifying psychological barriers to adherence. They can offer support by providing counseling, discussing the impact of mental health on medication adherence, and referring patients to mental health professionals if needed. Additionally, pharmacists can create a non-judgmental environment where patients feel comfortable discussing their mental health concerns, offering reassurance and emphasizing the importance of continued medication use for managing both physical and mental health. Addressing psychological barriers helps ensure patients adhere to their medication regimen and improve their overall well-being **(Horvat et al., 2024).**

The complexity of medication regimens is a significant barrier to adherence, particularly for patients managing multiple chronic conditions. Complex dosing schedules, intricate drug regimens, and medications with varying administration routes can overwhelm patients, leading to confusion, mistakes, and missed doses **(Talens et al., 2021).** Pharmacists can help by simplifying regimens where possible. They can collaborate with healthcare providers to reduce the number of medications or adjust the timing and frequency of doses to create a more straightforward plan. Pharmacists can also educate patients on the proper use of medications and the importance of following instructions accurately. Providing written instructions, visual aids, and reminders can help patients navigate their complex regimens. By making medication regimens simpler and easier to manage, pharmacists improve adherence rates and ensure better health outcomes for chronic disease patients **(Konstantinou et al., 2020).**

Social and environmental factors, including living conditions, lack of social support, or limited access to healthcare, can also affect medication adherence. Patients in low-income or rural areas may struggle with transportation, which can prevent them from accessing pharmacies or healthcare appointments. Additionally, some patients may not have family or community support to help them with medication management **(Alyami et al., 2024).** Pharmacists can help identify and address these barriers by offering solutions like home delivery services, providing information about local transportation options, or connecting patients with community resources. Pharmacists can also collaborate with nurses to involve caregivers in the treatment process, ensuring that patients have the necessary social support to adhere to their medications. By addressing these external factors, pharmacists ensure that patients can overcome environmental barriers and maintain consistent medication adherence **(Patton et al., 2021).**

The absence of regular follow-up or monitoring can contribute to poor medication adherence. Without consistent check-ins from healthcare providers, patients may feel disconnected from their treatment plan and lose motivation to continue taking their medications. Pharmacists can support adherence by providing ongoing monitoring and follow-up **(Bastani et al., 2021).** Through medication therapy management (MTM) services, pharmacists can track patients' medication usage, check for potential drug interactions, and address any emerging concerns. Regular follow-up phone calls or reminders can also help ensure patients are staying on track with their regimens. By maintaining consistent communication with patients and their healthcare teams, pharmacists play an essential role in identifying issues early on and preventing medication non-adherence from escalating into a more significant problem **(Alemede et al., 2024).**

Nurses are in a unique position to observe patients' adherence behaviors and identify potential barriers. They can serve as the first line of defense in recognizing when patients are struggling with their medication regimens. Pharmacists can support nurses by providing training and tools that help them identify signs of non-adherence, such as missed doses, incorrect usage, or refusal to take medications

**(Gutierrez & Sakulbumrungsil, 2021).** Nurses can then relay this information to the pharmacist, allowing for prompt intervention. Pharmacists can also work with nurses to educate patients about their medications, emphasizing the importance of adherence and addressing any concerns they may have. Together, pharmacists and nurses can create a supportive environment where patients are empowered to adhere to their medications and receive continuous care **(Hughes et al., 2022).**

Patient counseling is one of the most effective ways pharmacists can help improve medication adherence. By providing patients with clear, concise information about their medications, pharmacists can reduce confusion and empower patients to take ownership of their health. Counseling should include information about the medication's purpose, how it should be taken, potential side effects, and what to do in case of missed doses **(Yang et al., 2022).** Pharmacists should also encourage patients to ask questions, ensuring that they fully understand their treatment plan. In addition to one-on-one counseling, pharmacists can provide written materials, brochures, or access to online resources that patients can refer to later. This empowers patients to make informed decisions about their medication regimen and encourages consistent adherence, ultimately leading to better health outcomes **(Gatwood et al., 2022).**

Collaboration between pharmacists and nurses is essential to overcoming the barriers to medication adherence. Nurses often have direct, ongoing contact with patients, while pharmacists have the specialized knowledge to address medication-related issues **(Hamrick et al., 2020).** By working together, pharmacists and nurses can identify adherence barriers, share insights, and create individualized strategies to help patients stay on track with their medications. This collaboration can involve regular meetings to discuss patient progress, adjustments to treatment regimens, or joint patient education efforts. With a unified approach, pharmacists and nurses can provide patients with the support they need to overcome barriers to adherence, ensuring better outcomes for chronic disease management **(Ekong et al., 2020).**

Medication adherence in chronic disease patients requires ongoing, long-term support. Chronic conditions often require lifelong medication management, and patients need continued reinforcement to maintain adherence over time **(Okolo et al., 2024).** Pharmacists can provide long-term support through regular follow-up consultations, medication reviews, and adherence counseling. They can also help monitor any changes in the patient’s health that may affect their treatment, such as new diagnoses or changes in lifestyle. Nurses can complement this by offering routine check-ins, emotional support, and encouragement, especially for patients who may feel discouraged by the challenges of managing a chronic condition. By offering continuous support and fostering long-term relationships, both pharmacists and nurses help ensure that patients remain committed to their treatment regimens, improving their overall health and quality of life **(Alemede et al., 2024).**

**References**


1. Abdulhussein, N. N., Al-Azzawi, A. K., & Ajel, A. R. (2024, October). Design and implementation of closed-loop drug management in the pharmaceutical industry. In *AIP Conference Proceedings* (Vol. 3232, No. 1). AIP Publishing.
2. Al-Anzi, E. A. M., Al Harbi, Y. S. S., Al-Harbi, A. A. H. A., Al-Anzi, F. A. T., Alenazi, M. A. O., Alenazi, M. K. M., ... & Al Mutairi, A. Z. S. (2024). Publics’ Perceptions of Community Pharmacists role in Patient Care Services. *Journal of International Crisis and Risk Communication Research*, 1400-1407.
3. Al-Dosari, D. F., Al-Dosari, D. F., Al-Massifri, H. S. M., Alrajhi, E. A., Al-Jumaiah, N. S., Alenazi, R. H., & Al-Dosari, A. N. (2024). Enhancing Patient Outcomes: The Role of Pharmacists in OTC Drug Selection and Counseling. *Journal of International Crisis and Risk Communication Research*, 2038-2045.
4. Alemede, V., Nwankwo, E. I., Igwama, G. T., Olaboye, J. A., & Anyanwu, E. C. (2024). Pharmacists as educators: Enhancing patient understanding and access to specialty medications through community workshops. *Magna Scientia Advanced Biology and Pharmacy*, *13*(1), 1-9.
5. Alemede, V., Nwankwo, E. I., Igwama, G. T., Olaboye, J. A., & Anyanwu, E. C. (2024). Designing state-level policies to support independent pharmacies in providing specialty care services in rural regions. *Magna Scientia Advanced Biology and Pharmacy*, *13*(1), 19-29.

6. Al-Jaroodi, J., Mohamed, N., & Abukhousa, E. (2020). Health 4.0: on the way to realizing the healthcare of the future. Ieee Access, 8, 211189-211210.
7. Alqarni, M. A., Alshammari, N. A., Aljohani, A. A., Alenezi, B. H., Alsubhi, I. M., Al-Harbi, H. R., ... & Aljabry, S. (2023). The Role Of Community Pharmacists In Educating Patients About Drug Interactions And Medication Safety. *Journal of Survey in Fisheries Sciences*, *10*(5), 276-281.
8. Alsaihati, F. M., Alsultan, F. M., Bujubarah, M. M., Alsharyawi, F. M. S., Aljaffar, B. H., & Hussain, F. (2024). Integrative Approaches in Primary Care: The Roles of Dentistry, Nursing, and Pharmacy. *Letters in High Energy Physics,643-656*
9. Al-Worafi, Y. M. (2023). Pharmaceutical Care. In *Handbook of Medical and Health Sciences in Developing Countries: Education, Practice, and Research* (pp. 1-23). Cham: Springer International Publishing.
10. Alyami, A. M. O., Alsulayyi, M. S. M., Alruwaili, F. N. S., Alruwaili, A. S. A., Alenezi, F. S. K., Alruwaili, R. M. M., ... & Najai, A. M. N. (2024). Role of Nurses in Improving Medication Adherence among Elderly Patients. *Journal of International Crisis and Risk Communication Research*, 310-332.
11. Balogun, O. D., Ayo-Farai, O., Ogundairo, O., Maduka, C. P., Okongwu, C. C., Babarinde, A. O., & Sodamade, O. T. (2023). Innovations in drug delivery systems: A review of the pharmacist's role in enhancing efficacy and patient compliance. *World Journal of Advanced Research and Reviews*, *20*(3), 1268-1282.
12. Balogun, O. D., Ayo-Farai, O., Ogundairo, O., Maduka, C. P., Okongwu, C. C., Babarinde, A. O., & Sodamade, O. T. (2024). The role of pharmacists in personalised medicine: a review of integrating pharmacogenomics into clinical practice. *International Medical Science Research Journal*, *4*(1), 19-36.
13. Bastani, P., Bikineh, P., Ravangard, R., Rezaee, R., & Kavosi, Z. (2021). Determinants affecting medication adherence in the elderly: A qualitative study. *Aging Medicine*, *4*(1), 35-41.
14. Bethishou, L. (2024). The Role of Pharmacy in Supporting Safe and Effective Transitions of Care. *Journal of Contemporary Pharmacy Practice*, *71*(2), 23-33.
15. Bhattarai, B., Walpola, R., Khan, S., & Mey, A. (2021). Factors associated with medication adherence among people living with COPD: Pharmacists' perspectives. *Exploratory Research in Clinical and Social Pharmacy*, *3*, 100049.
16. Burnier, M. (2024). The role of adherence in patients with chronic diseases. *European Journal of Internal Medicine*, *119*, 1-5.
17. Chavan, A. A., Kumbhar, S. B., Shinde, V. R., Thorat, A. A., Jadhav, P. B., Jadhav, A. R., ... & Kumbhar, D. D. (2023). Role of pharmacist in healthcare system. *GSC Biological and Pharmaceutical Sciences*, *24*(1).
18. Coman, L. I., Ianculescu, M., Paraschiv, E. A., Alexandru, A., & Bădărău, I. A. (2024). Smart Solutions for Diet-Related Disease Management: Connected Care, Remote Health Monitoring Systems, and Integrated Insights for Advanced Evaluation. *Applied Sciences*, *14*(6), 2351.
19. Coutsouvelis, J., Siderov, J., Tey, A. Y., Bortz, H. D., O'Connor, S. R., Rowan, G. D., ... & Percival, M. A. (2020). The impact of pharmacist-led strategies implemented to reduce errors related to cancer therapies: a systematic review. *Journal of Pharmacy Practice and Research*, *50*(6), 466-480.
20. Darlow, B., Brown, M., Hudson, B., Frew, G., Clark, J., Vincent, L., ... & Briggs, A. M. (2023). Knee osteoarthritis and the knowledgeable, trustworthy pharmacist: Patient and pharmacist perceptions of community pharmacy-based education and support. *Musculoskeletal Care*, *21*(1), 3-15.
21. De Baetselier, E., Dilles, T., Batalha, L. M., Dijkstra, N. E., Fernandes, M. I., Filov, I., ... & Van Rompaey, B. (2021). Perspectives of nurses' role in interprofessional pharmaceutical care across 14 European countries: A qualitative study in pharmacists, physicians and nurses. *PloS one*, *16*(5), e0251982.
22. Dilles, T., Heczkova, J., Tziaferi, S., Helgesen, A. K., Grøndahl, V. A., Van Rompaey, B., ... & Jordan, S. (2021). Nurses and pharmaceutical care: interprofessional, evidence-based working to improve patient care and outcomes. *International journal of environmental research and public health*, *18*(11), 5973.
23. Ekong, G., Chou, C., Lakin, J., Hardin, A., Fox, B., Hunt, C., & Kavookjian, J. (2020). Pharmacist-led motivational interviewing for diabetes medication adherence in a worksite wellness program. *Journal of the American Pharmacists Association*, *60*(6), e224-e229.

24. Ferreri, S. P., Hughes, T. D., & Snyder, M. E. (2020). Medication therapy management: current challenges. *Integrated Pharmacy Research and Practice*, 71-81.
25. Forsyth, P., Radley, A., Rushworth, G. F., Marra, F., Roberts, S., O'Hare, R., ... & Maguire, B. (2023). The collaborative care model: realizing healthcare values and increasing responsiveness in the pharmacy workforce. *Research in social and administrative pharmacy*, *19*(1), 110-122.
26. Gatwood, J., Brooks, C., Meacham, R., Abou-Rahma, J., Cernasev, A., Brown, E., & Kuchtey, R. W. (2022). Facilitators and barriers to glaucoma medication adherence. *Journal of Glaucoma*, *31*(1), 31-36.
27. Gil-Guillen, V. F., Balsa, A., Bernárdez, B., Valdés y Llorca, C., Márquez-Contreras, E., de la Haba-Rodríguez, J., ... & Gómez-Martínez, J. (2022). Medication non-adherence in rheumatology, oncology and cardiology: a review of the literature of risk factors and potential interventions. *International Journal of Environmental Research and Public Health*, *19*(19), 12036.
28. Gutierrez, M. M., & Sakulbumrungsil, R. (2021). Factors associated with medication adherence of hypertensive patients in the Philippines: a systematic review. *Clinical hypertension*, *27*, 1-15.
29. Haleem, A., Javaid, M., Singh, R. P., & Suman, R. (2021). Telemedicine for healthcare: Capabilities, features, barriers, and applications. *Sensors international*, *2*, 100117.
30. Hamrahian, S. M., Maarouf, O. H., & Fülöp, T. (2022). A critical review of medication adherence in hypertension: barriers and facilitators clinicians should consider. *Patient preference and adherence*, 2749-2757.
31. Hamrick, J., Augustine, J. M., Pinkerton, D., Klein, C. M., & Elliott, J. (2020). Simulated medication adherence: First and second-year pharmacy students' perceptions of adhering to a multi-drug medication regimen. *Currents in Pharmacy Teaching and Learning*, *12*(7), 858-863.
32. Horvat, M., Eržen, I., & Vrbnjak, D. (2024, August). Barriers and facilitators to medication adherence among the vulnerable elderly: a focus group study. In *Healthcare* (Vol. 12, No. 17, p. 1723). MDPI.
33. Hughes, T. D., Cutrell, S., Minshew, L. M., Brown, P., & Ferreri, S. P. (2022). An exploration of barriers, facilitators, and practical solutions for adopting medication synchronization into community pharmacies: A qualitative analysis. *Exploratory Research in Clinical and Social Pharmacy*, *5*, 100111.
34. Jaffer, U., Nassir, C. M. N. C. M., & Ahmed, M. A. (2023). Medication adherence: understanding the challenges among the elderly. *Psychology*, *8*(52), 865-874.
35. Jarab, A. S., Al-Qerem, W., Mukattash, T. L., Abuhishmah, S. R., & Alkhdour, S. (2022). Pharmacists' knowledge and attitudes toward medication therapy management service and the associated challenges and barriers for its implementation. *Saudi Pharmaceutical Journal*, *30*(6), 842-848.
36. Kasahun, A. E., Sendekie, A. K., Mekonnen, G. A., Sema, F. D., Kemal, L. K., & Abebe, R. B. (2022). Impact of personal, cultural and religious beliefs on medication adherence among patients with chronic diseases at university hospital in Northwest Ethiopia. *Patient preference and adherence*, 1787-1803.
37. Kelley, A. T., Incze, M. A., Baylis, J. D., Calder, S. G., Weiner, S. J., Zickmund, S. L., ... & Bridges, J. F. (2022). Patient-centered quality measurement for opioid use disorder: Development of a taxonomy to address gaps in research and practice. *Substance abuse*, *43*(1), 1286-1299.
38. Kiles, T. M., Peroulas, D., & Borja-Hart, N. (2022). Defining the role of pharmacists in addressing the social determinants of health. *Research in Social and Administrative Pharmacy*, *18*(9), 3699-3703.
39. Konstantinou, P., Kassianos, A. P., Georgiou, G., Panayides, A., Papageorgiou, A., Almas, I., ... & Karekla, M. (2020). Barriers, facilitators, and interventions for medication adherence across chronic conditions with the highest non-adherence rates: a scoping review with recommendations for intervention development. *Translational behavioral medicine*, *10*(6), 1390-1398.
40. Kvarnström, K., Westerholm, A., Airaksinen, M., & Liira, H. (2021). Factors contributing to medication adherence in patients with a chronic condition: a scoping review of qualitative research. *Pharmaceutics*, *13*(7), 1100.
41. Lott, B. E., Anderson, E. J., Zapata, L. V., Cooley, J., Forbes, S., Taylor, A. M., ... & Warholak, T. (2021). Expanding pharmacists' roles: Pharmacists' perspectives on barriers and facilitators to collaborative practice. *Journal of the American Pharmacists Association*, *61*(2), 213-220.

42. Ngamdee, N., Chumfang, N., Chiangkhong, A., Satasuwan, A., & Sukdee, S. (2024). Behavioral Patterns and Barriers to Medication Adherence in Older Adults with Diabetes. *International Journal of Geoinformatics*, *20*(3), 64-73.
43. O'Toole, J., Krishnan, M., Riekert, K., & Eakin, M. N. (2022). Understanding barriers to and strategies for medication adherence in COPD: a qualitative study. *BMC pulmonary medicine*, *22*(1), 98.
44. Oakley, C., & Ream, E. (2024, February). Role of the Nurse in Patient Education and Engagement and Its Importance in Advanced Breast Cancer. In *Seminars in Oncology Nursing* (Vol. 40, No. 1, p. 151556). WB Saunders.
45. Okolo, C. A., Arowoogun, J. O., Chidi, R., & Adeniyi, A. O. (2024). Telemedicine's role in transforming healthcare delivery in the pharmaceutical industry: A systematic review. *World Journal of Advanced Research and Reviews*, *21*(2), 1836-1856.
46. Passey, D. G., Healy, R., Qualls, J., Halwani, A., & Sauer, B. C. (2021). Pharmacist-led collaborative medication management programs for oral antineoplastic therapies: A systematic literature review. *Journal of the American Pharmacists Association*, *61*(3), e7-e18.
47. Patton, D. E., Ryan, C., & Hughes, C. M. (2021). Enhancing community pharmacists' provision of medication adherence support to older adults: a mixed methods study using the theoretical domains framework. *Research in social and administrative pharmacy*, *17*(2), 406-418.
48. Pruette, C. S., & Amaral, S. (2021). Empowering patients to adhere to their treatment regimens: a multifaceted approach. *Pediatric transplantation*, *25*(1), e13849.
49. Rahayu, S. A., Widianto, S., Defi, I. R., & Abdulah, R. (2021). Role of pharmacists in the interprofessional care team for patients with chronic diseases. *Journal of multidisciplinary healthcare*, 1701-1710.
50. Ravi, P., Pfaff, K., Ralph, J., Cruz, E., Bellaire, M., & Fontanin, G. (2022). Nurse-pharmacist collaborations for promoting medication safety among community-dwelling adults: A scoping review. *International Journal of Nursing Studies Advances*, *4*, 100079.
51. Raza, M. A., Aziz, S., Noreen, M., Anjum, I., & Raza, S. M. (2022). A portrait of the pharmacy profession globally: pharmacist universal professional identity and establishment of global pharmacy council. *INNOVATIONS in pharmacy*, *13*(1).
52. Schmid, S., Schlosser, S., Gülow, K., Pavel, V., Müller, M., & Kratzer, A. (2022). Interprofessional collaboration between ICU physicians, staff nurses, and hospital pharmacists optimizes antimicrobial treatment and improves quality of care and economic outcome. *Antibiotics*, *11*(3), 381.
53. Schneider, M. P., & Burnier, M. (2023). Partnership between patients and interprofessional healthcare providers along the multifaceted journey to medication adherence. *British Journal of Clinical Pharmacology*, *89*(7).
54. Seixas, A., Conners, C., Chung, A., Donley, T., & Jean-Louis, G. (2020). A pantheoretical framework to optimize adherence to healthy lifestyle behaviors and medication adherence: the use of personalized approaches to overcome barriers and optimize facilitators to achieve adherence. *JMIR mHealth and uHealth*, *8*(6), e16429.
55. Sugandh, F. N. U., Chandio, M., Raveena, F. N. U., Kumar, L., Karishma, F. N. U., Khuwaja, S., & Kumar, S. (2023). Advances in the management of diabetes mellitus: a focus on personalized medicine. *Cureus*, *15*(8).
56. Syrnyk, M., & Glass, B. (2023). Pharmacist interventions in medication adherence in patients with mental health disorders: a scoping review. *International Journal of Pharmacy Practice*, *31*(5), 449-458.
57. Tadros, E., Barbini, M., & Kaur, L. (2023). Collaborative healthcare in incarcerated settings. *International Journal of Offender Therapy and Comparative Criminology*, *67*(9), 910-929.
58. Talens, A., Guilabert, M., Lumbreras, B., Aznar, M. T., & López-Pintor, E. (2021). Medication experience and adherence to oral chemotherapy: A qualitative study of patients' and health professionals' perspectives. *International Journal of Environmental Research and Public Health*, *18*(8), 4266.
59. Toscos, T., Drouin, M., Pater, J. A., Flanagan, M., Wagner, S., Coupe, A., ... & Mirro, M. J. (2020). Medication adherence for atrial fibrillation patients: triangulating measures from a smart pill bottle,

e-prescribing software, and patient communication through the electronic health record. *JAMIA open*, *3*(2), 233-242.

60. Wheat, L., Roane, T. E., Connelly, A., Zeigler, M., Wallace, J., Kim, J. H., & Segal, R. (2020). Using a pharmacist–community health worker collaboration to address medication adherence barriers. *Journal of the American Pharmacists Association*, *60*(6), 1009-1014.
61. While, A. (2020). Medication adherence: understanding the issues and finding solutions. *British Journal of Community Nursing*, *25*(10), 474-479.
62. White, A., Fulda, K. G., Blythe, R., Chui, M. A., Reeve, E., Young, R., ... & Xiao, Y. (2022). Defining and enhancing collaboration between community pharmacists and primary care providers to improve medication safety. *Expert opinion on drug safety*, *21*(11), 1357-1364.
63. Yang, C., Lee, D. T. F., Wang, X., & Chair, S. Y. (2022). Effects of a nurse-led medication self-management intervention on medication adherence and health outcomes in older people with multimorbidity: a randomised controlled trial. *International Journal of Nursing Studies*, *134*, 104314.